

## ON THE TAYLOR COEFFICIENTS OF RATIONAL FUNCTIONS

BY K. MAHLER

Communicated by J. W. S. CASSELS

*Received 22 February 1955; and in revised form 18 April 1955*

Dedicated to C. L. SIEGEL

Let  $F(z)$  be a rational function of  $z$  which is regular at  $z = 0$  and so possesses a convergent power series

$$F(z) = \sum_{h=0}^{\infty} f_h z^h.$$

The problem arises of characterizing those rational functions  $F(z)$  that have *infinitely many vanishing Taylor coefficients*  $f_h$ . After earlier and more special results by Siegel (2) and Ward (4) I applied in 1934 (1) a  $p$ -adic method due to Skolem (3) to the problem and obtained the following partial solution.

**THEOREM 1.** *Assume that all Taylor coefficients  $f_h$  of the rational function  $F(z)$  are algebraic numbers, and that infinitely many of them vanish. Then two integers  $L$  and  $L_1$  with  $0 \leq L_1 < L$  exist such that  $f_h$  is zero for all sufficiently large  $h \equiv L_1 \pmod{L}$ .*

In the present paper, the restriction on the character of the coefficients  $f_h$  will be removed, by showing the

**THEOREM 2.** *Theorem 1 remains valid when the coefficients  $f_h$  of  $F(z)$  are arbitrary complex numbers.*

In the proof of this theorem, the assertion will be reduced to one relating to rational functions with algebraic Taylor coefficients, and it will be assumed that the truth of Theorem 1 has already been established.

1. If the difference of two functions is a polynomial, all but finitely many of their Taylor coefficients are the same. Also to a given rational function one can always add a unique polynomial such that the sum function vanishes at the point at infinity.

Hence, without loss of generality, we shall assume from now on that the rational function  $F(z)$  is not only regular at  $z = 0$ , but also it vanishes at  $z = \infty$ . We then call  $F(z)$  a *normed function*. The restriction to normed functions considerably shortens the discussion.

2. Let  $L$  and  $L_1$  be two integers such that  $0 \leq L_1 < L$ . We say that  $F(z)$  has the *zero sequence*  $L_1 \pmod{L}$  if all but finitely many of the Taylor coefficients  $f_h$  with  $h \equiv L_1 \pmod{L}$  are zero.

This property may also be expressed in another form. Put

$$\epsilon = e^{2\pi i/L} \quad \text{and} \quad E(z) = \sum_{j=0}^{L-1} \epsilon^{jL_1} F(\epsilon^{-j}z).$$

Evidently 
$$E(z) = \sum_{h=0}^{\infty} f_h z^h \sum_{j=0}^{L-1} \epsilon^{j(L_1-h)} = L \sum_{\substack{h=0 \\ h \equiv L_1 \pmod{L}}}^{\infty} f_h z^h,$$

and so  $E(z)$  reduces to a polynomial if  $L_1 \pmod{L}$  is a zero sequence of  $F(z)$ . On the other hand, as  $F(z)$  is normed, all terms  $\epsilon^{jL_1} F(\epsilon^{-j}z)$  of  $E(z)$  vanish at  $z = \infty$ . The same is then true for  $E(z)$  itself, and so  $E(z)$  *vanishes identically*. Hence the stronger property

$$f_h = 0 \text{ for all suffixes } h \equiv L_1 \pmod{L}$$

holds.

3. Assume again that  $L_1 \pmod{L}$  is a zero sequence of  $F(z)$ . Let further  $\alpha_1, \alpha_2, \dots, \alpha_n$  be the distinct poles of  $F(z)$ ; by hypothesis, none of these poles lies at  $z = 0$ . Then  $F(\epsilon^{-j}z)$  has the poles

$$\epsilon^j \alpha_1, \quad \epsilon^j \alpha_2, \quad \dots, \quad \epsilon^j \alpha_n.$$

As was shown in § 2,

$$E(z) = \sum_{j=0}^{L-1} \epsilon^{jL_1} F(\epsilon^{-j}z) \equiv 0.$$

Hence the poles of  $F(z)$  are cancelled by the poles of the  $L-1$  other functions  $\epsilon^{jL_1} F(\epsilon^{-j}z)$  where  $j = 1, 2, \dots, L-1$ .

It follows therefore that *to every pole  $\alpha_\nu$  of  $F(z)$  there is a second pole  $\alpha_\mu$  ( $\mu \neq \nu$ ) such that  $\alpha_\nu / \alpha_\mu \neq 1$  is an  $L$ -th root of unity*, which, of course, need not be primitive. Furthermore,  $F(z)$  *has at least two distinct poles*.

4. Let  $\Sigma = \{\alpha_\nu / \alpha_\mu\}$  be the set of all those quotients  $\alpha_\nu / \alpha_\mu \neq 1$  of distinct poles of  $F(z)$  that are roots of unity. Unless  $\Sigma$  is the null set, there exists a smallest positive integer  $M$  such that  $\Sigma$  consists only of  $M$ th roots of unity which, however, need not all be primitive.

Assume, in particular, that  $L_1 \pmod{L}$  is a zero sequence of  $F(z)$ , and put

$$(L, M) = L^*, \quad L' = \frac{L}{L^*},$$

so that

$$L^* = L\Lambda + MM, \quad L = L^*L',$$

with certain integers  $\Lambda$  and  $M$ . By § 3,  $\Sigma$  is now certainly not the null set, because it contains elements that are  $L$ th roots of unity. Denote by  $\Sigma^*$  the subset formed by all these elements of  $\Sigma$  that are  $L$ th roots of unity. Thus the elements  $\alpha_\nu / \alpha_\mu$  of  $\Sigma^*$  satisfy both equations

$$\left(\frac{\alpha_\nu}{\alpha_\mu}\right)^L = 1 \quad \text{and} \quad \left(\frac{\alpha_\nu}{\alpha_\mu}\right)^M = 1,$$

and so also the equation

$$\left(\frac{\alpha_\nu}{\alpha_\mu}\right)^{L^*} = \left\{\left(\frac{\alpha_\nu}{\alpha_\mu}\right)^L\right\}^\Lambda \left\{\left(\frac{\alpha_\nu}{\alpha_\mu}\right)^M\right\}^M = 1.$$

Therefore  $\Sigma^*$  *consists only of  $L^*$ th roots of unity*.

5. We introduce now the  $L'$  new functions

$$E_k(x) = \sum_{\substack{j=0 \\ j \equiv k \pmod{L'}}}^{L-1} \epsilon^{jL_1} F(\epsilon^{-j}z) \quad (k = 0, 1, 2, \dots, L' - 1).$$

As already shown, the sum of these functions

$$E(z) = \sum_{k=0}^{L'-1} E_k(z) = \sum_{j=0}^{L-1} \epsilon^{jL_1} F(\epsilon^{-j}z)$$

is *identically zero*.

It is obvious that  $E_0(z)$  may have poles only at the points  $\epsilon^j\alpha_1, \epsilon^j\alpha_2, \dots, \epsilon^j\alpha_n$ , where  $j \equiv 0 \pmod{L'}$ ,  $0 \leq j \leq L-1$ , while, for  $k = 1, 2, \dots, L'-1$ , poles of  $E_k(z)$  can lie only at  $\epsilon^t\alpha_1, \epsilon^t\alpha_2, \dots, \epsilon^t\alpha_n$ , where  $t \equiv k \pmod{L'}$ ,  $0 \leq t \leq L-1$ . Let us suppose that  $\epsilon^j\alpha_\nu$  is a pole of  $E_0(z)$ , and that  $\epsilon^t\alpha_\mu$  is one of  $E_k(z)$ , where  $1 \leq k \leq L'-1$ . Then

$$\iota - j \equiv k \not\equiv 0 \pmod{L'}.$$

Hence  $L^*(\iota - j) \not\equiv 0 \pmod{L}$ ,

whence  $(\epsilon^{\iota-j})^{L^*} \neq 1, \quad \epsilon^{\iota-j} \neq 1$ .

Therefore, necessarily,

$$\epsilon^j\alpha_\nu \neq \epsilon^t\alpha_\mu,$$

because,  $\epsilon^{\iota-j}$  being an  $L$ th root of unity, the quotient

$$\frac{\alpha_\nu}{\alpha_\mu} = \epsilon^{\iota-j} \neq 1$$

would otherwise belong to  $\Sigma^*$  and be an  $L^*$ th root of unity; and this is not the case.

The function  $E_0(z)$  has then no poles in common with the other terms  $E_k(z)$  of  $E(z)$ , and all its poles are also poles of  $E(z)$ . Since  $E(z)$  has no poles,  $E_0(z)$  is thus a polynomial. But, from its definition in terms of  $F(z)$ ,  $E_0(z)$  is a *normed* rational function. Hence, finally,

*$E_0(z)$  is identically zero.*

Put now

$$\eta = \epsilon^{L'} = e^{2\pi i/L^*}.$$

Evidently  $E_0(z) = \sum_{j=0}^{L^*-1} \eta^{jL_1} F(\eta^{-j}z) = L^* \sum_{\substack{h=0 \\ h \equiv L_1 \pmod{L^*}}}^{\infty} f_h z^h \equiv 0,$

whence  $f_h = 0$  for all suffixes  $h \equiv L_1 \pmod{L^*}$ .

The following result has thus been established.

LEMMA 1. *Let  $L_1 \pmod{L}$  be a zero sequence of  $F(z)$ ; let  $\alpha_1, \alpha_2, \dots, \alpha_n$  be the distinct poles of  $F(z)$ ; and let  $M$  be the smallest integer such that all quotients  $\frac{\alpha_\mu}{\alpha_\nu} \neq 1$ , that are roots of unity, are  $M$ -th roots of unity. If  $L^* = (L, M)$ , then  $F(z)$  admits the zero sequence  $L_1 \pmod{L^*}$ .*

This lemma is of importance for later, because  $L^*$  is a divisor of  $M$ , and  $M$  depends only on the poles of  $F(z)$ . We note that the lemma remains valid when  $F(z)$  is not normed, but shall not use this fact.

6. We proceed now to the proof of Theorem 2.

The most general rational function  $F(z) \not\equiv 0$  regular at  $z = 0$  is of the form

$$F(z) = \frac{a_0 + a_1 z + \dots + a_m z^m}{(z - \alpha_1)^{e_1} (z - \alpha_2)^{e_2} \dots (z - \alpha_n)^{e_n}}.$$

Here  $e_1, e_2, \dots, e_n$  are arbitrary positive integers;  $a_0, a_1, \dots, a_m$  are arbitrary complex numbers with  $a_m \neq 0$ ; and  $\alpha_1, \alpha_2, \dots, \alpha_n$ , the poles of  $F(z)$ , are complex numbers that are all distinct and different from zero, but are otherwise arbitrary. The function  $F(z)$  is assumed to be normed, and therefore the inequality

$$m < e_1 + e_2 + \dots + e_n$$

holds.

If again

$$F(z) = \sum_{h=0}^{\infty} f_h z^h$$

is the power series of  $F(z)$  in the neighbourhood of  $z = 0$ , let  $H$  be the set of all suffixes  $h$  for which

$$f_h = 0.$$

It is assumed that  $H$  is an infinite set; the problem is to prove that under this hypothesis  $F(z)$  possesses at least one zero sequence.

7. From now on let  $X$  be the set of the  $m + n + 1$  parameters

$$X = \{a_0, a_1, \dots, a_m, \alpha_1^{-1}, \alpha_2^{-1}, \dots, \alpha_n^{-1}\}$$

that occur in  $F(z)$ ; the use of  $\alpha_v^{-1}$  rather than  $\alpha_v$  will prove to be an advantage. Further put

$$e_0 = e_1 + e_2 + \dots + e_n.$$

Then  $F(z)$  may also be written in the form

$$F(z) = (-1)^{e_0} \prod_{\nu=1}^n (\alpha_\nu^{-1})^{e_\nu} \sum_{\mu=0}^m a_\mu z^\mu \prod_{\nu=1}^n (1 - \alpha_\nu^{-1} z)^{-e_\nu}.$$

On developing here the last factor into a power series by means of the binomial theorem, we see immediately that, for  $h = 0, 1, 2, \dots$ ,  $f_h$  is a polynomial with rational coefficients in the elements of  $X$ .

Hence, if  $X$  consists only of algebraic numbers, the coefficients  $f_h$  are likewise algebraic. It is assumed that this is no longer the case; hence  $X$  includes at least one transcendental number.

Denote by  $R$  the Gaussian imaginary quadratic field. The elements of  $X$  generate a smallest extension field

$$P = R(X) = R(a_0, a_1, \dots, a_m, \alpha_1^{-1}, \alpha_2^{-1}, \dots, \alpha_n^{-1})$$

over  $R$ . It is shown in algebra that this extension field may be constructed as follows.

We first adjoin to  $R$  a certain finite set of transcendental complex numbers

$$\sigma_1, \sigma_2, \dots, \sigma_p$$

that are algebraically independent over  $R$ , so arriving at the purely transcendental extension

$$P_0 = R(\sigma_1, \sigma_2, \dots, \sigma_p).$$

The field  $P$  is now derived from  $P_0$  by a simple algebraic extension

$$P = P_0(\tau) = R(\sigma_1, \sigma_2, \dots, \sigma_p, \tau),$$

$\tau$  being a suitable complex number algebraic over  $P_0$ .

This number  $\tau$  may still be chosen in many different ways, and there is no loss of generality in assuming that  $\tau$  is *integral over the polynomial ring*  $R[\sigma_1, \sigma_2, \dots, \sigma_p]$ . The equation for  $\tau$  has then the form

$$Q(\sigma_1, \sigma_2, \dots, \sigma_p; \tau) \equiv \tau^q + \sum_{\kappa=1}^q Q_\kappa(\sigma_1, \sigma_2, \dots, \sigma_p) \tau^{q-\kappa} = 0,$$

where 
$$Q_\kappa(\sigma_1, \sigma_2, \dots, \sigma_p) \quad (\kappa = 1, 2, \dots, q)$$

are polynomials in  $R[\sigma_1, \sigma_2, \dots, \sigma_p]$ . It may also be assumed that  $Q(\sigma_1, \sigma_2, \dots, \sigma_p; \tau)$ , considered as a polynomial in  $\sigma_1, \sigma_2, \dots, \sigma_p, \tau$ , is *irreducible over*  $R$ .

8. The elements  $a_\mu$  and  $\alpha_\nu^{-1}$  of  $X$  are finite numbers in  $P$ . They can therefore be written as polynomials in  $\tau$ , with coefficients that are rational functions of  $\sigma_1, \sigma_2, \dots, \sigma_p$  with numerical coefficients in  $R$ . Denote by  $\Delta(\sigma_1, \sigma_2, \dots, \sigma_p) \neq 0$  the least common denominator of these rational functions;  $\Delta$  is thus an element of  $R[\sigma_1, \sigma_2, \dots, \sigma_p]$ . Then  $a_\mu$  and  $\alpha_\nu^{-1}$  take the form

$$a_\mu = \frac{A_\mu(\sigma_1, \sigma_2, \dots, \sigma_p; \tau)}{\Delta(\sigma_1, \sigma_2, \dots, \sigma_p)} \quad (\mu = 0, 1, \dots, m)$$

and 
$$\alpha_\nu^{-1} = \frac{\Lambda_\nu(\sigma_1, \sigma_2, \dots, \sigma_p; \tau)}{\Delta(\sigma_1, \sigma_2, \dots, \sigma_p)} \quad (\nu = 1, 2, \dots, n).$$

Here the numerators

$$A_\mu(\sigma_1, \sigma_2, \dots, \sigma_p; \tau), \quad \Lambda_\nu(\sigma_1, \sigma_2, \dots, \sigma_p; \tau)$$

belong to the polynomial ring  $R[\sigma_1, \sigma_2, \dots, \sigma_p, \tau]$ .

On substituting these expressions for the elements of  $X$ ,  $F(z)$  becomes a rational function

$$F(z) = \Phi(z \mid \sigma_1, \sigma_2, \dots, \sigma_p; \tau)$$

not only of  $z$ , but also of  $\sigma_1, \sigma_2, \dots, \sigma_p, \tau$ , while its numerical coefficients lie in  $R$ . It follows further, from the representation of  $f_h$  as a polynomial in the elements of  $X$  with coefficients in  $R$ , that these Taylor coefficients may be written as

$$f_h = \frac{\phi_h(\sigma_1, \sigma_2, \dots, \sigma_p; \tau)}{\Delta(\sigma_1, \sigma_2, \dots, \sigma_p)^{d_h}} \quad (h = 0, 1, 2, \dots),$$

where the numerators 
$$\phi_h(\sigma_1, \sigma_2, \dots, \sigma_p; \tau)$$

lie in the polynomial ring  $R[\sigma_1, \sigma_2, \dots, \sigma_p, \tau]$ , while the exponents  $d_h$  are certain positive integers depending on  $h$ . One may, in fact, choose  $d_h = e_0 + h + 1$ ; but we shall not need this. The hypothesis on  $f_h$  implies that

$$\phi_h(\sigma_1, \sigma_2, \dots, \sigma_p; \tau) = 0 \quad \text{if} \quad h \in H.$$

9. Let us now replace the algebraically independent complex numbers  $\sigma_1, \sigma_2, \dots, \sigma_p$  by independent complex variables

$$s_1, s_2, \dots, s_p,$$

and the complex number  $\tau$  for which

$$Q(\sigma_1, \sigma_2, \dots, \sigma_p; \tau) = 0$$

by a dependent complex variable  $t$  satisfying

$$Q(s_1, s_2, \dots, s_p; t) = 0.$$

We then obtain a new rational function

$$F^*(z) = \Phi(z \mid s_1, s_2, \dots, s_p; t)$$

of  $z$ , as well as of  $s_1, s_2, \dots, s_p, t$ , with numerical coefficients in  $R$ . This function has the explicit form

$$F^*(z) = \frac{a_0^* + a_1^* z + \dots + a_m^* z^m}{(z - \alpha_1^*)^{e_1} (z - \alpha_2^*)^{e_2} \dots (z - \alpha_n^*)^{e_n}},$$

where

$$a_\mu^* = \frac{A_\mu(s_1, s_2, \dots, s_p; t)}{\Delta(s_1, s_2, \dots, s_p)} \quad (\mu = 0, 1, \dots, m)$$

and

$$\alpha_\nu^* = \frac{A_\nu(s_1, s_2, \dots, s_p; t)}{\Delta(s_1, s_2, \dots, s_p)} \quad (\nu = 1, 2, \dots, n).$$

Further it possesses the power series

$$F^*(z) = \sum_{h=0}^{\infty} f_h^* z^h,$$

where

$$f_h^* = \frac{\phi_h(s_1, s_2, \dots, s_p; t)}{\Delta(s_1, s_2, \dots, s_p)^{d_h}} \quad (h = 0, 1, 2, \dots).$$

Since  $\Delta$  does not vanish identically, and since the change from  $\sigma_1, \sigma_2, \dots, \sigma_p, \tau$  to  $s_1, s_2, \dots, s_p, t$  maps  $P = R(\sigma_1, \sigma_2, \dots, \sigma_p, \tau)$  isomorphically onto  $R(s_1, s_2, \dots, s_p, t)$ , it is clear that also

$$\phi_h(s_1, s_2, \dots, s_p, t) = 0 \quad \text{and} \quad f_h^* = 0 \quad \text{if} \quad h \in H.$$

It is further obvious from the construction that for

$$s_1 = \sigma_1, \quad s_2 = \sigma_2, \quad \dots, \quad s_p = \sigma_p, \quad t = \tau,$$

the equations

$$F^*(z) = F(z), \quad a_\mu^* = a_\mu, \quad \alpha_\nu^* = \alpha_\nu, \quad f_h^* = f_h$$

hold.

10. To simplify the notation, we introduce the  $p$ -dimensional space  $C^p$  of all points

$$\mathbf{s} = (s_1, s_2, \dots, s_p), \quad \mathbf{s}' = (s'_1, s'_2, \dots, s'_p), \quad \boldsymbol{\sigma} = (\sigma_1, \sigma_2, \dots, \sigma_p), \quad \dots,$$

with arbitrary real or complex coordinates, and we make  $C^p$  a metric space by defining the distance  $\rho(\mathbf{s}, \mathbf{s}')$  of two points  $\mathbf{s}, \mathbf{s}'$  by

$$\rho(\mathbf{s}, \mathbf{s}') = \{ |s_1 - s'_1|^2 + |s_2 - s'_2|^2 + \dots + |s_p - s'_p|^2 \}^{\frac{1}{2}}.$$

Let  $R^p$  similarly be the set of all points in  $C^p$  the coordinates of which lie in the Gaussian field  $R$ ; thus  $R^p$  is dense in  $C^p$ . We can then select in many ways an infinite sequence of points

$$S = \{\mathbf{s}^{(1)}, \mathbf{s}^{(2)}, \mathbf{s}^{(3)}, \dots\}, \quad \text{where} \quad \mathbf{s}^{(k)} = (s_1^{(k)}, s_2^{(k)}, \dots, s_p^{(k)}),$$

in  $R^p$  such that

$$\lim_{k \rightarrow \infty} \mathbf{s}^{(k)} = \boldsymbol{\sigma}, \quad \text{i.e.} \quad \lim_{k \rightarrow \infty} \rho(\mathbf{s}^{(k)}, \boldsymbol{\sigma}) = 0.$$

From the form of the equation

$$Q(s_1, s_2, \dots, s_p, t) = 0$$

for  $t$ , it is further possible to associate with each point  $\mathbf{s}^{(k)}$  of  $S$  a complex root,  $t^{(k)}$  say, of the equation

$$Q(s_1^{(k)}, s_2^{(k)}, \dots, s_p^{(k)}, t^{(k)}) = 0,$$

such that also

$$\lim_{k \rightarrow \infty} t^{(k)} = \tau.$$

11. Denote, for  $k = 1, 2, 3, \dots$ , by

$$F^{(k)}(z), \quad a_\mu^{(k)}, \quad \alpha_\nu^{(k)}, \quad f_h^{(k)},$$

the expressions into which  $F^*(z), \quad a_\mu^*, \quad \alpha_\nu^*, \quad f_h^*$ ,

respectively, are changed on putting

$$s_1 = s_1^{(k)}, \quad s_2 = s_2^{(k)}, \quad \dots, \quad s_p = s_p^{(k)}, \quad t = t^{(k)}.$$

Then  $F^{(k)}(z)$  is the rational function

$$F^{(k)}(z) = \frac{\alpha_0^{(k)} + \alpha_1^{(k)}z + \dots + \alpha_m^{(k)}z^m}{(z - \alpha_1^{(k)})^{e_1} (z - \alpha_2^{(k)})^{e_2} \dots (z - \alpha_n^{(k)})^{e_n}} = \Phi(z \mid s_1^{(k)}, s_2^{(k)}, \dots, s_p^{(k)}, t^{(k)})$$

of  $z$  with the Taylor series

$$F^{(k)}(z) = \sum_{h=0}^{\infty} f_h^{(k)} z^h,$$

and here

$$f_h^{(k)} = 0 \quad \text{if} \quad h \in H.$$

We must, however, assume that  $k$  is already sufficiently large, i.e. that  $\mathbf{s}^{(k)}$  is sufficiently near to  $\boldsymbol{\sigma}$ , so as to exclude the possibility that one of the expressions  $a_\mu^{(k)}, \alpha_\nu^{(k)}, f_h^{(k)}$  becomes infinite, or that one of the poles  $\alpha_\nu^{(k)}$  vanishes, or that two of these poles coincide. Assume, say, that these cases are excluded when  $k \geq k_0$ .

It follows now, from the continuity properties of a rational function, that

$$\lim_{k \rightarrow \infty} a_\mu^{(k)} = a_\mu, \quad \lim_{k \rightarrow \infty} \alpha_\nu^{(k)} = \alpha_\nu, \quad \lim_{k \rightarrow \infty} f_h^{(k)} = f_h$$

for all values of the suffixes  $\mu, \nu$  and  $h$ .

12. The equation  $Q(s_1^{(k)}, s_2^{(k)}, \dots, s_p^{(k)}, t^{(k)}) = 0$

for  $t^{(k)}$  is of degree  $q$ , and its coefficients lie in  $R$ ; for both the numerical coefficients of  $Q$ , and the coordinates of  $\mathbf{s}^{(k)}$ , belong to  $R$ . Therefore  $t^{(k)}$  is an algebraic number at most of degree  $q$  over the Gaussian field, hence at most of degree  $2q$  over the rational field. Denote by  $K^{(k)} = R(t^{(k)})$  the algebraic extension of  $R$  generated by  $t^{(k)}$ ; this field has likewise a degree not greater than  $2q$  over the rational field. From their definitions, it is clear that the numbers

$$a_\mu^{(k)}, \quad \alpha_\nu^{(k)}, \quad f_h^{(k)}$$

all are elements of  $K^{(k)}$ , as soon as  $k \geq k_0$ .

In particular, the Taylor coefficients  $f_h^*$  of

$$F^{(k)}(z) = \sum_{h=0}^{\infty} f_h^{(k)} z^h$$

are algebraic numbers, and furthermore infinitely many of these coefficients vanish,

$$f_h^{(k)} = 0 \quad \text{if} \quad h \in H.$$

The hypothesis of Theorem 1 is thus satisfied. Hence, for every  $k \geq k_0$ ,  $F^{(k)}(z)$  possesses at least one zero sequence  $L_1 \pmod{L}$ . Here we may assume that  $0 \leq L_1 < L$ . Both  $L = L^{(k)}$  and  $L_1 = L_1^{(k)}$  may still depend on  $k$ . We note that, by hypothesis,

$$m < e_1 + e_2 + \dots + e_n.$$

Hence also  $F^{(k)}(z)$  is normed, so that *all* its Taylor coefficients  $f_h^{(k)}$  satisfying  $h \equiv L_1 \pmod{L}$  are zero.

13. Lemma 1 enables us to construct a zero sequence  $L_1 \pmod{L}$  of  $F^{(k)}(z)$  with *bounded*  $L$ , hence also with *bounded*  $L_1$ .

The poles  $\alpha_1^{(k)}, \alpha_2^{(k)}, \dots, \alpha_n^{(k)}$  of  $F^{(k)}(z)$  lie in  $K^{(k)}$ , and the same is therefore true for the quotients of two such poles. Denote by  $\Sigma = \Sigma^{(k)}$  the set of all those quotients

$$\frac{\alpha_\mu^{(k)}}{\alpha_\nu^{(k)}} \neq 1$$

that are roots of unity; we know, from § 3, that  $\Sigma$  is not the null set. Hence a smallest positive integer  $M = M^{(k)}$  exists such that all elements of  $\Sigma$  are  $M$ th roots of unity.

By Lemma 1,  $F^{(k)}(z)$  admits also the larger zero sequence

$$L_1 \pmod{L^*}, \quad \text{where } L^* = (L, M).$$

This zero sequence is identical with  $L_1^* \pmod{L^*}$ , where  $L_1^*$  is the integer for which

$$L_1^* \equiv L_1 \pmod{L^*}, \quad 0 \leq L_1^* < L^*.$$

The roots of unity which are the elements of  $\Sigma$  lie in the algebraic field  $K^{(k)}$ , and this field is at most of degree  $2q$ . On the other hand, there are only finitely many roots of unity that are algebraic numbers at most of degree  $2q$ . Denote by  $M_0$  the least common multiple of the orders of all these roots of unity. Then evidently

$$M^{(k)} \leq M_0 \quad \text{for } k \geq k_0.$$

Since  $L^*$  is a divisor of  $M^{(k)}$ , this implies that also

$$0 \leq L_1^* < L^* \leq M_0 \quad \text{for } k \geq k_0.$$

14. On dropping now again the asterisk, the last result may be formulated as follows:

*If  $k \geq k_0$ , then  $F^{(k)}(z)$  possesses at least one zero sequence*

$$L_1 \pmod{L}, \quad \text{where } 0 \leq L_1 < L \leq M_0,$$

*and where  $M_0$  is independent of  $k$ . Moreover, all Taylor coefficients  $f_h^{(k)}$  of  $F^{(k)}(z)$  with  $h \equiv L_1 \pmod{L}$  are zero.*

There exist only *finitely many* zero sequences  $L_1 \pmod{L}$  for which  $0 \leq L_1 < L \leq M_0$ , the zero sequences

$$Z_1, \quad Z_2, \quad \dots, \quad Z_v,$$

say. For each  $k \geq k_0$  denote by  $u = u^{(k)}$  the smallest suffix such that  $Z_u$  is a zero sequence of  $F^{(k)}(z)$ . This function  $u = u^{(k)}$  has only  $v$  possible values. Hence there is an infinite sequence of indices

$$\dot{k} = k_1, k_2, k_3, \dots, \quad \text{where } k_0 \leq k_1 \leq k_2 \leq k_3 \leq \dots,$$



for which  $u = u^{(k)}$  assumes one and the same fixed value  $u_0$ . For all these indices  $F^{(k)}(z)$  possesses the same zero sequence  $Z_{u_0}$ , or say  $L_1^0 \pmod{L^0}$ , and all Taylor coefficients  $f_h^{(k)}$  with  $h \equiv L_1^0 \pmod{L^0}$  are zero. However, as was proved in § 11,

$$\lim_{k \rightarrow \infty} f_h^{(k)} = f_h \quad \text{for all } h.$$

Therefore, on allowing  $k$  to run over the sequence  $k_1, k_2, k_3, \dots$  to infinity, it follows at once that also

$$f_h = 0 \quad \text{if } h \equiv L_1^0 \pmod{L^0}.$$

Hence the original function  $F(z)$  likewise admits the zero sequence  $L_1^0 \pmod{L^0}$ . This proves the assertion.

15. Theorem 2 implies a slightly stronger result.

THEOREM 3. Let 
$$F(z) = \sum_{h=0}^{\infty} f_h z^h$$

be a rational function of  $z$  which is regular at  $z = 0$  and has infinitely many vanishing Taylor coefficients  $f_h$ . Then a positive integer  $L$  and at most  $L$  non-negative integers  $L_1, L_2, \dots, L_l$  with

$$L_j \not\equiv L_k \pmod{L} \quad \text{for } j \neq k$$

exist such that  $f_h$  vanishes exactly when

$$h \equiv L_j \pmod{L}, \quad h \geq L_j \quad (j = 1, 2, \dots, l)$$

and for at most finitely many other values of  $h$ .

*Proof.* It may again be assumed that  $F(z)$  is normed. Denote by  $M$  the same positive integer as in Lemma 1. By this lemma, it suffices to consider those zero sequences  $L_j \pmod{L}$  of  $F(z)$  for which  $L$  is a divisor of  $M$ . As such sequences can be subdivided into sequences  $\pmod{M}$ , it further suffices to prove the theorem with  $L$  replaced by  $M$ .

Denote by  $L_1, L_2, \dots, L_l \pmod{M}$  all distinct residue classes  $h \equiv L_j \pmod{M}$  that contain infinitely many suffixes  $h$  for which  $f_h = 0$ . The assertion is proved if it can be shown that each  $L_j \pmod{M}$  is a zero sequence of  $F(z)$ . It will be enough to consider  $L_1 \pmod{M}$ .

We assume thus that

$$f_h = 0 \quad \text{for infinitely many } h \equiv L_1 \pmod{M}.$$

Similarly as before, put

$$e = e^{2\pi i/M}, \quad E(z) = \sum_{j=0}^{M-1} e^{L_1 j} F(e^{-j} z);$$

further write

$$\zeta = z^M.$$

Then 
$$z^{-L_1} E(z) = M z^{-L_1} \sum_{\substack{h=0 \\ h \equiv L_1 \pmod{M}}}^{\infty} f_h z^h = M \sum_{k=0}^{\infty} f_{L_1+kM} z^{kM} = E(\zeta),$$

where

$$E(\zeta) = M \sum_{k=0}^{\infty} f_{L_1+kM} \zeta^k$$

evidently is a rational function of  $\zeta$ . This new function  $E(\zeta)$  is regular at  $\zeta = 0$ , vanishes at  $\zeta = \infty$ , and has infinitely many vanishing Taylor coefficients  $f_{L_1+kM}$ .

Hence it follows from Theorem 2 that  $E(\zeta)$  possesses at least one zero sequence

$$k \equiv \kappa_1 \pmod{\kappa}.$$

As the function is normed, this implies that

$$f_{L_1+kM} = 0 \quad \text{if} \quad k \equiv \kappa_1 \pmod{\kappa},$$

or, what is the same,

$$f_h = 0 \quad \text{if} \quad h \equiv L_1 + \kappa_1 M \pmod{\kappa M}.$$

This relation means that the original function  $F(z)$  has the zero sequence  $L_1 + \kappa_1 M \pmod{\kappa M}$ . But then, by Lemma 1, it also admits the larger zero sequence  $L_1 + \kappa_1 M \pmod{M}$ , hence also the zero sequence  $L_1 \pmod{M}$ . This concludes the proof.

16. It is well known that, for sufficiently large  $h$ , the Taylor coefficient  $f_h$  of the rational function  $F(z)$  has the explicit representation

$$f_h = \sum_{\nu=1}^n p_\nu(h) \beta_\nu^h$$

where  $p_1(h), p_2(h), \dots, p_n(h)$  are polynomials in the variable  $h$  not identically zero, while  $\beta_1, \beta_2, \dots, \beta_n$  are distinct constants different from zero, viz. the reciprocals of the poles of  $F(z)$ . Conversely, every expression of this kind defines the Taylor coefficients of a rational function regular at  $z = 0$ , and the same is true if  $h$  is replaced by  $-h$ . The following result is then implicit in Theorem 2.

**THEOREM 4.** *Let  $\beta_1, \beta_2, \dots, \beta_n$  be finitely many complex numbers that are distinct, different from zero, and such that no quotient of two of them is a root of unity. Also let  $p_1(h), p_2(h), \dots, p_n(h)$  be an equal number of polynomials not identically zero with arbitrary complex coefficients. Then the equation*

$$\sum_{\nu=1}^n p_\nu(h) \beta_\nu^h = 0$$

*has at most finitely many solutions in rational integers.*

I am very much indebted to my colleague, Dr P. Cohn, for reading the manuscript and suggesting many improvements, and to the referee, Dr T. Estermann, for a great simplification of the proof.

#### REFERENCES

- (1) MAHLER, K. *Proc. Acad. Sci. Amst.* 38 (1935), 51–60.
- (2) SIEGEL, C. L. *Tôhoku math. J.* 20 (1921), 26–31.
- (3) SKOLEM, TH. *Skr. norske Vidensk. Akad.* (1933), no. 6.
- (4) WARD, M. *Math. Z.* 39 (1934), 211–14.