

MONTY

SO, UH, WHY IS YOUR NAME "MR. PI"? DO YOU LIKE PIES OR SOMETHING?

THE PRIVY COUNCIL ON RIGEL-9 GAVE ME THIS NAME BEFORE EXILING ME FROM THE PLANET...



www.comics.com

MOST RIGELIANS HAVE RATIONAL NUMBERS FOR NAMES LIKE "14" OR "286733", BUT BECAUSE I'M HALF HUMAN AND TAINTED WITH HUMAN BLOOD THEY CHOSE TO LABEL ME WITH THE IRRATIONAL NUMBER "PI".



6/8

OK. WAIT. WAIT. WAIT.



©2004 Jim Meddick / Dist. by NEA, Inc.

"PI" IS A NUMBER?



I THOUGHT HUMANS WERE REQUIRED TO LEARN MATHEMATICS IN SCHOOL.




SOME DO. SOME BUY CALCULATORS.





(BEEP)
HELLO, YOU'VE
REACHED
THE FOX
RESIDENCE.



TO LEAVE A MESSAGE,
PRESS THE SQUARE ROOT
OF 1,296 MINUS THE CUBE
ROOT OF 13,824 TIMES
17.5 MINUS THE 4TH ROOT
OF 1,908,029,761.



SORRY,
TIME'S UP.
GOODBYE.
(CLICK)



I'VE FIGURED
OUT WHAT'S
THE MATTER.
WITH THE
ANSWERING
MACHINE,
OR OUR
SON?

PSYCHIATRIC
CLINIC

TAKE A
NUMBER


27



"TAKE A
NUMBER"?---
BUT I
HAVE
MATH
ANXIETY!

Mathematics by Experiment:

Plausible Reasoning in the 21st Century



Jonathan M. Borwein, FRSC



Research Chair in IT
Dalhousie University

Halifax, Nova Scotia, Canada



ICME Regular Lecture

ICME10, Copenhagen, July 4–11, 2004

If mathematics describes an objective world just like physics, there is no reason why inductive methods should not be applied in mathematics just the same as in physics.

(Kurt Gödel, 1951)


URL: www.expmath.info




MY INTENTIONS IN THIS TALK

- to discuss Experimental Methodology*
 - its philosophy
 - its practice
 - using concrete amusing (I hope) examples

Insight taking place



Molecular Borromean Rings



- and then to
- review its past
 - describe its present
 - predict its future

- Thereby, to persuade you that traditional accounting of mathematical learning and research is largely an ahistorical caricature[†]


*Bailey, Moll and I invite you to the 2nd [Experimental Mathematics Workshop](#) at Tulane, Spring 2005

[†]Online is a superset of this talk

FROM TROTSKY to GÖDEL: a **LIFE**

"By 1948, the Marxist-Leninist ideas about the proletariat and its political capacity seemed more and more to me to disagree with reality ... I pondered my doubts, and for several years the study of mathematics was all that allowed me to preserve my inner equilibrium. Bolshevik ideology was, for me, in ruins. I had to build another life."

Jean Van Heijenoort (1913-1986) *With Trotsky in Exile*, in Anita Feferman's *From Trotsky to Gödel*



THE EVIL of BOURBAKI

“There is a story told of the mathematician Claude Chevalley (1909–84), who, as a true Bourbaki, was extremely opposed to the use of images in geometric reasoning.



He is said to have been giving a very abstract and algebraic lecture when he got stuck. After a moment of pondering, he turned to the blackboard, and, trying to hide what he was doing, drew a little diagram, looked at it for a moment, then quickly erased it, and turned back to the audience and proceeded with the lecture. ...

... The computer offers those less expert, and less stubborn than Chevalley, access to the kinds of images that could only be imagined in the heads of the most gifted mathematicians, ... ” (Nathalie Sinclair^a)

^aChapter in *Making the Connection: Research and Practice in Undergraduate Mathematics*, MAA Notes, 2004 in Press.

Abstract. I shall talk generally about experimental and heuristic mathematics, and give accessible, primarily visual and symbolic, examples.

The emergence of powerful mathematical computing environments like *Maple* and *Matlab*, the growing availability of correspondingly powerful (*multi-processor*) computers and the pervasive presence of the internet allow for research mathematicians, students and teachers, to proceed heuristically and ‘quasi-inductively’.

We may increasingly use symbolic and numeric computation visualization tools, simulation and data mining. Likewise, an aesthetic appreciation of mathematics may be provided to a much broader audience.

Many of the benefits of computation are accessible through low-end 'electronic blackboard' versions of experimental mathematics. This also permits livelier classes, more realistic examples, and more collaborative learning. Moreover, the distinction between computing (HPC) and communicating (HPN) is increasingly moot.

The unique features of the discipline make this both more problematic and more challenging.

- For example, there is still no truly satisfactory way of displaying mathematical notation on the web;
- and we care more about the reliability of our literature than does any other science.

The traditional role of proof in mathematics is arguably under siege.

Limited by examples, I intend to ask:

- ★ What constitutes secure mathematical knowledge?
- ★ When is computation convincing? Are humans less fallible?
 - What tools are available? What methodologies?
 - What about the 'law of the small numbers'?
 - Who cares for certainty? What is the role of proof?
- ★ How is mathematics actually done? How should it be?

And I shall offer some personal conclusions.

- ▶ Many of the more sophisticated examples originate in the boundary between mathematical physics and number theory and involve the ζ -function, $\zeta(n) = \sum_{k=1}^{\infty} \frac{1}{k^n}$, and its relatives.

They often rely on the sophisticated use of *Integer Relations Algorithms* — recently ranked among the ‘top ten’ algorithms of the century.

- **Integer Relation methods** were first discovered by our colleague **Helaman Ferguson** the mathematical sculptor.

See www.cecm.sfu.ca/projects/IntegerRelations/

SCIENCE NEWS April 23 2004

www.cs.dal.ca/~jborwein

MATH

A Digital Slice of Pi


THE NEW WAY TO DO PURE MATH: EXPERIMENTALLY BY W. WAYT GIBBS

“One of the greatest ironies of the information technology revolution is that while the computer was conceived and born in the field of pure mathematics, through the genius of giants such as John von Neumann and Alan Turing, until recently this marvelous technology had only a minor impact within the field that gave it birth.” So begins *Experimentation in Mathematics*, a book by Jonathan M. Borwein and David H. Bailey due out in September that documents how all that has begun to change. Computers, once looked on by mathematical researchers with disdain as mere calculators, have gained enough power to enable an entirely new way to make fundamental discoveries: by running experiments and observing what happens.

The first clear evidence of this shift emerged in 1996. Bailey, who is chief technologist at the National Energy Research Sci-

entific Computing Center in Berkeley, Calif., and several colleagues developed a computer program that could uncover integer relations among long chains of real numbers. It was a problem that had long vexed mathematicians. Euclid discovered the first integer relation scheme—a way to work out the greatest common divisor of any two integers—around 300 B.C. But it wasn't until 1977 that Helaman Ferguson and Rodney W. Forcade at last found a method to detect relations among an arbitrarily large set of numbers. Building on that work, in 1995 Bailey's group turned its computers loose on some of the fundamental constants of math, such as log 2 and pi.

To the researchers' great surprise, after months of calculations the machines came up with novel formulas for these and other nat-



COMPUTER RENDERINGS

of mathematical constructs can reveal hidden structure. The bands of color that appear in this plot of all solutions to a certain class of polynomials [specifically, those of the form $\pm 1 \pm x \pm x^2 \pm x^3 \pm \dots \pm x^n = 0$, up to $n = 18$] have yet to be explained by conventional analysis.

Computer experiments are transforming mathematics

BY ERICA KLARREICH

Many people regard mathematics as the crown jewel of the sciences. Yet math has historically lacked one of the defining trappings of science: laboratory equipment. Physicists have their particle accelerators; biologists, their electron microscopes; and astronomers, their telescopes. Mathematics, by contrast, concerns not the physical landscape but an idealized, abstract world. For exploring that world, mathematicians have traditionally had only their intuition.

Now, computers are starting to give mathematicians the lab instrument that they have been missing. Sophisticated software is enabling researchers to travel further and deeper into the mathematical universe. They're calculating the number pi with mind-boggling precision, for instance, or discovering patterns in the contours of beautiful, infinite chains of spheres that arise out of the geometry of knots.

Experiments in the computer lab are leading mathematicians to discoveries and insights that they might never have reached by traditional means. "Pretty much every [mathematical] field has been transformed by it," says Richard Crandall, a mathematician at Reed College in Portland, Ore. "Instead of just being a number-crunching tool, the computer is becoming more like a garden shovel that turns over rocks, and you find things underneath."

At the same time, the new work is raising unsettling questions about how to regard experimental results in fields for which

"I have some of the excitement that Leonardo of Pisa must have felt when he encountered Arabic arithmetic. It suddenly made certain calculations flabbergastingly easy," Borwein says. "That's what I think is happening with computer experimentation today."

EXPERIMENTERS OF OLD In one sense, math experiments are nothing new. Despite their field's reputation as a purely deductive science, the great mathematicians over the centuries have never limited themselves to formal reasoning and proof.


For instance, in 1666, sheer curiosity and love of numbers led Isaac Newton to calculate directly the first 16 digits of the number pi, later writing, "I am ashamed to tell you to how many figures I carried these computations, having no other business at the time."

Carl Friedrich Gauss, one of the towering figures of 19th-century mathematics, habitually discovered new mathematical results by experimenting with numbers and looking for patterns. When Gauss was a teenager, for instance, his experiments led him to one of the most important conjectures in the history of number theory: that the number of prime numbers less than a number x is roughly equal to x divided by the logarithm of x .


Gauss often discovered results experimentally long before he could prove them formally. Once, he complained, "I have the result, but I do not yet know how to get it."

In the case of the prime number theorem, Gauss later refined his conjecture but never did figure out how to prove it. It took more than a century for mathematicians to come up with a proof.

Like today's mathematicians, math experimenters in the late 19th century used computers—but in those days, the word referred to people with a special facility for calculating



UNSOLVED MYSTERIES — A computer experiment produced this plot of all the solutions to a collection of simple equations in 2001. Mathematicians are still trying to account for its many features.



STRAIGHT CIRCLES — When mathematicians Colin Adams and Eric Schoenfeld created this image while playing with the computer program Snappea last year, they were stunned to see perfectly straight chains of spheres. The observation led them to an unexpected discovery about knots.

Mathematicians, Bailey and Borwein argue, should be free to work more like other scientists do, developing hypotheses through experimentation and then testing them in further experiments. Formal proof is still the ideal, they say, but it is not the only path to mathematical truth.

“When I started school, I thought mathematics was about proofs, but now I think it’s about having secure mathematical knowledge,” Borwein says. “We claim that’s not the same thing.”

Bailey and Borwein point out that mathematical proofs can run to hundreds of pages and require such specialized knowledge that only a few people are capable of reading and judging them.

“We feel that in many cases, computations constitute very strong evidence, evidence that is at least as compelling as some of the more complex formal proofs in the literature,” Bailey and Borwein say in *Mathematics by Experiment*.

“One thing that’s happening is you can discover many more things than you can explain.”

— JONATHAN BORWEIN
DALHOUSIE UNIVERSITY



mathematics,

n. **1.** a group of related subjects, including algebra, geometry, trigonometry and calculus, concerned with the study of number, quantity, shape, and space, and their inter-relationships, applications, generalizations and abstractions. It is not too much of a parody of the history of mathematics to see it as developing from the arithmetical description of a few common-place concerns, albeit that the results obtained at this early stage, by, for example, Babylonian and Chinese astronomers, were both sophisticated and accurate. As this process of arithmetization expanded, more general conceptions of number were developed, together with the crucial insight that these results had general validity, and represented functional relationships; the lack of suitable notations for variables and functions, however, hampered the development of mathematics for a millennium. Although it was recognized at an early stage that justification requires rigorous proof which must ultimately rest upon unexceptionable axioms (until the limitations of the axiomatic method were proven in the present century), and for two millennia Euclid's treatment of geometry was regarded as the paradigm not only of mathematics but of science in general, the present concept of rigor in proof and definition, as exemplified by epsilon-delta notation, largely originates with Cauchy and others in the 19th century. The search for certainty also gave rise, as in Descartes' analytic geometry, to a search for

FINDING THINGS or PROVING THINGS



Consider the following two *Euler sum identities* both discovered heuristically.

- Both merit quite firm belief—more so than many proofs.

Why?

- Only the first warrants significant effort for its proof.

Why and **Why Not?**



I. A MULTIPLE ZETA VALUE

Euler sums or *MZVs* are a wonderful generalization of the classical ζ function.

For natural numbers

$$\zeta(i_1, i_2, \dots, i_k) := \sum_{n_1 > n_2 > \dots > n_k > 0} \frac{1}{n_1^{i_1} n_2^{i_2} \dots n_k^{i_k}}$$

◇ Thus $\zeta(a) = \sum_{n \geq 1} n^{-a}$ is as before and

$$\zeta(a, b) = \sum_{n=1}^{\infty} \frac{1 + \frac{1}{2^b} + \dots + \frac{1}{(n-1)^b}}{n^a}$$

✓ k is the sum's *depth* and $i_1 + i_2 + \dots + i_k$ is its *weight*.

- This clearly extends to alternating and character sums.

- MZV's satisfy many striking identities, of which the simplest are

$$\zeta(2, 1) = \zeta(3) \quad 4\zeta(3, 1) = \zeta(4).$$

- MZV's have recently found interesting interpretations in high energy physics, knot theory, combinatorics ...
- ✓ Euler found and partially proved theorems on **reducibility** of depth 2 to depth 1 ζ 's
- $\zeta(6, 2)$ is the lowest weight **'irreducible'**.
- ✓ High precision *fast ζ -convolution* (see *EZ-Face/Java*) allows use of integer relation methods and leads to important dimensional (reducibility) conjectures and amazing identities.

A STRIKING CONJECTURE open for all $n > 2$ is:

$$8^n \zeta(\{-2, 1\}_n) \stackrel{?}{=} \zeta(\{2, 1\}_n),$$

There is abundant evidence amassed since it was found in 1996.* For example, very recently Petr Lisonek checked the first 85 cases to 1000 places in about 41 HP hours with only the *expected error*. And $N=163$ in ten hours.

- This is the *only* identification of its type of an Euler sum with a distinct MZV.
- Can even just the case $n = 2$ be proven *symbolically* as is the case for $n = 1$?

*Equivalently that the functions

$$L_{-2,1}(1, 2t) = L_{2,1}(1, t) \quad (= L_3(1, t)),$$

defined later agree for small t .

II. A CHARACTER EULER SUM

Let

$$[2b, -3](s, t) := \sum_{n>m>0} \frac{(-1)^{n-1}}{n^s} \frac{\chi_3(m)}{m^t},$$

where χ_3 is the character modulo 3. Then for $N = 0, 1, 2, \dots$

$$\begin{aligned} & [2b, -3](2N + 1, 1) \\ &= \frac{L_{-3}(2N + 2)}{4^{1+N}} - \frac{1 + 4^{-N}}{2} L_{-3}(2N + 1) \log(3) \\ &+ \sum_{k=1}^N \frac{1 - 3^{-2N+2k}}{2} L_{-3}(2N - 2k + 2) \alpha(2k) \\ &- \sum_{k=1}^N \frac{1 - 9^{-k}}{1 - 4^{-k}} \frac{1 + 4^{-N+k}}{2} L_{-3}(2N - 2k + 1) \alpha(2k + 1) \\ &- 2L_{-3}(1) \alpha(2N + 1). \end{aligned}$$

✓ Here α is the *alternating zeta function* and L_{-3} is the *primitive L-series modulo 3*.

★ One first **evaluates** such sums as integrals

DICTIONARIES ARE LIKE TIMEPIECES

- ▶ Samuel Johnson observed of watches that “the best do not run true, and the worst are better than none.” The same is true of tables and databases. Michael Berry “would give up Shakespeare in favor of Prudnikov, Brychkov and Marichev.”

- That excellent compendium contains

$$(1) \quad \sum_{k=1}^{\infty} \sum_{l=1}^{\infty} \frac{1}{k^2 (k^2 - kl + l^2)} = \frac{\pi^{\alpha} \sqrt{3}}{30},$$


where the “ α ” is *probably* “4” [volume 1, entry 9, page 750].

- ★ Integer relation methods suggest that **no reasonable value of α works.**

What is intended in (1)?

POLYA and HEURISTICS

[I]ntuition comes to us much earlier and with much less outside influence than formal arguments which we cannot really understand unless we have reached a relatively high level of logical experience and sophistication.



In the first place, the beginner must be convinced that proofs deserve to be studied, that they have a purpose, that they are interesting.

(George Polya*, 1968)

*In *Mathematical Discovery: On Understanding, Learning and Teaching Problem Solving*.

MOORE'S LAW

The complexity for minimum component costs has increased at a rate of roughly a factor of two per year. ... Certainly over the short term this rate can be expected to continue, if not to increase. Over the longer term, the rate of increase is a bit more uncertain, although there is no reason to believe it will not remain nearly constant for at least 10 years.

(Gordon Moore, Intel co-founder, 1965)

► “Moore’s Law” asserts that semiconductor technology approximately doubles in capacity and performance roughly every 18 to 24 months (not quite every year as Moore predicted).

This trend has continued unabated for 40 years, and, according to Moore and others, there is still no end in sight—at least another ten years is assured.

► This astounding record of sustained exponential progress has no peer in the history of technology.



What's more, mathematical computing tools are now being implemented on parallel computer platforms, which will provide even greater power to the research mathematician.

► Amassing huge amounts of processing power will not solve all mathematical problems, even those amenable to computational analysis.

There are cases where a dramatic increase in computation could, by itself, result in significant breakthroughs, but it is easier to find examples where this is unlikely to happen.

SIMON and RUSSELL

This skyhook-skyscraper construction of science from the roof down to the yet unconstructed foundations was possible because the behaviour of the system at each level depended only on a very approximate, simplified, abstracted characterization at the level beneath.¹³

This is lucky, else the safety of bridges and airplanes might depend on the correctness of the “Eightfold Way” of looking at elementary particles.

- ◇ Herbert A. Simon, *The Sciences of the Artificial*, MIT Press, 1996, page 16. (An early experimental computational scientist.)

13... More than fifty years ago Bertrand Russell made the same point about the architecture of mathematics. See the "Preface" to *Principia Mathematica* "... the chief reason in favour of any theory on the principles of mathematics must always be inductive, i.e., it must lie in the fact that the theory in question allows us to deduce ordinary mathematics. In mathematics, the greatest degree of self-evidence is usually not to be found quite at the beginning, but at some later point; hence the early deductions, until they reach this point, give reason rather for believing the premises because true consequences follow from them, than for believing the consequences because they follow from the premises." Contemporary preferences for deductive formalisms frequently blind us to this important fact, which is no less true today than it was in 1910.

VISUAL DYNAMICS


- In recent continued fraction work, we needed to study the *dynamical system* $t_0 := t_1 := 1$:

$$t_n \leftrightarrow \frac{1}{n} t_{n-1} + \omega_{n-1} \left(1 - \frac{1}{n}\right) t_{n-2},$$

where $\omega_n = a^2, b^2$ for n even, odd respectively.


✓ Think of this as a **black box**.

- ▷ Numerically all one sees is $t_n \rightarrow 0$ slowly.
- ▷ Pictorially we *learn* significantly more*:



*... “Then felt I like a watcher of the skies, when a new planet swims into his ken.” (*Chapman’s Homer*)

- Scaling by \sqrt{n} , and coloring **odd** and **even** iterates, **fine structure appears**.




The **attractors** for various $|a| = |b| = 1$.

- ★ This is now fully explained with a *lot* of work—the rate of convergence in some cases by a fine *singular-value* argument.

GAUSS and HADAMARD

Carl Friedrich Gauss, who drew (carefully) and computed a great deal, once noted, *I have the result, but I do not yet know how to get it.**



Pauca sed Matura

The object of mathematical rigor is to sanction and legitimize the conquests of intuition, and there was never any other object for it.

- ◇ J. Hadamard quoted at length in E. Borel, *Lecons sur la theorie des fonctions*, 1928.

*Likewise the quote!

De lemniscata, elegantissime omnes exactissimas
 ones superantia acquisitiones et quibus
 per methodos quae campum propositum
 neminem nobis aperuunt. Gott. Jul.

Solutio problematis ballistici Gott. Jul.

Comentarum theoriam perfectivam reddidi Gott. Jul.

Novus in analysi campus se nobis aperuit,
 scilicet investigatio functionum etc.

Formas superiores considerare coepimus
 Mr. Febr. 1798

Formulas exactas pro parallaxi
 eruimus Mr. Apr. 8.

Terminum medium arithmetico-geometricum
 inter 1 et $\sqrt{2}$ esse = $\frac{\pi}{10}$ vixque
 ad hujusmodi videremur comprobavimus, quare
 demonstrata proinde novus campus in analysi
 certis aperietur Mr. Mai 30.

In principis Geometriae egregios propositus
 terminus Mr. Sept.

Circa terminos medios arithmetico-geometricos
 multae nova determinavit. Mr. Novemb.

Novus in analysi campus se nobis aperuit

An excited young Gauss writes: "A new field of analysis has appeared to us, evidently in the study of functions etc." (October 1798)


MOTIVATION and GOALS

INSIGHT – demands speed \equiv **micro-parallelism**

- For rapid verification.
- For validation; proofs *and* refutations; “monster barring” .
- ★ What is “easy” changes: HPC & HPN blur, merging disciplines and collaborators — democratizing mathematics but challenging authenticity.
- **Parallelism** \equiv more space, speed & stuff.
- **Exact** \equiv hybrid \equiv symbolic ‘+’ numeric (*Maple meets NAG*).
- In analysis, algebra, geometry & topology.

... MOREOVER

- Towards an Experimental **Methodology** — philosophy and practice.
- ▶ **Intuition is acquired** — mesh computation and mathematics.
- **Visualization** — 3 is a lot of dimensions.
- ▶ “Monster-barring” (Lakatos) and “Caging”:
 - randomized checks: equations, linear algebra, primality
 - graphic checks: compare $2\sqrt{y} - 2y$ and




EXPERIMENTAL MATHODOLOGY.

1. Gaining insight and intuition.
2. Discovering new patterns and relationships.
3. Graphing to expose math principles.
4. Testing and especially falsifying conjectures.
5. Exploring a possible result to see if it *merits* formal proof.
6. Suggesting approaches for formal proof.
7. Computing replacing lengthy hand derivations.
8. Confirming analytically derived results.

A BRIEF HISTORY OF RIGOUR

- Greeks: trisection, circle squaring, cube doubling and $\sqrt{2}$.
- Newton and Leibniz: fluxions and infinitesimals.
- Cauchy and Fourier: limits and continuity.
- Frege and Russell, Gödel and Turing.



Fourier series need not converge

THE PHILOSOPHIES OF RIGOUR

- Everyman: **Platonism**—stuff exists (1936)
- Hilbert: **Formalism**—math is invented; formal symbolic games without meaning
- Brouwer: **Intuitionism**—many variants; (embodied cognition)
- Bishop: **Constructivism**—tell me how big; (social constructivism)

† Last two deny the *excluded middle*: $A \vee \tilde{A}$

HALES and KEPLER

- Kepler's conjecture: **the densest way to stack spheres is in a pyramid** is the oldest problem in discrete geometry.
- The most interesting recent example of computer assisted proof. Published in *Annals of Math* with an “only 99% checked” disclaimer.
- This has triggered very varied reactions. (In Math, Computers Don't Lie. Or Do They? *NYT* 6/4/04)
- Famous earlier examples: **The Four Color Theorem** and **The non existence of a projective plane of order 10.**
- The three raise and answer quite distinct questions—both real and specious.

Does the proof stack up?

Think peer review takes too long? One mathematician has waited four years to have his paper refereed, only to hear that the exhausted reviewers can't be certain whether his proof is correct. George Szpiro investigates.



Grocers the world over know the most efficient way to stack spheres — but a mathematical proof for the method has brought reviewers to their knees.

Just under five years ago, Thomas Hales made a startling claim. In an e-mail he sent to dozens of mathematicians, Hales declared that he had used a series of computers to prove an idea that has evaded certain confirmation for 400 years. The subject of his message was Kepler's conjecture, proposed by the German astronomer Johannes Kepler, which states that the densest arrangement of spheres is one in which they are stacked in a pyramid — much the same way as grocers arrange oranges.

Soon after Hales made his announcement, reports of the breakthrough appeared on the front pages of newspapers around the world. But today, Hales's proof remains in limbo. It has been submitted to the prestigious *Annals of Mathematics*, but is yet to appear in print. Those charged with checking it say that they believe the proof is correct, but are so exhausted with the verification process that they cannot definitively rule out any errors. So when Hales's manuscript finally does appear in the *Annals*, probably during the next year, it will carry an unusual editorial note — a statement that parts of the paper have proved impossible to check.

At the heart of this bizarre tale is the use of computers in mathematics, an issue that has split the field. Sometimes described as a 'brute force' approach, computer-aided

proofs often involve calculating thousands of possible outcomes to a problem in order to produce the final solution. Many mathematicians dislike this method, arguing that it is inelegant. Others criticize it for not offering any insight into the problem under consideration. In 1977, for example, a computer-aided proof was published for the four-colour theorem, which states that no more than four colours are needed to fill in a map so that any two adjacent regions have different colours¹². No errors have been found in the proof, but some mathematicians continue to seek a solution using conventional methods.

Pile-driver

Hales, who started his proof at the University of Michigan in Ann Arbor before moving to the University of Pittsburgh, Pennsylvania, began by reducing the infinite number of possible stacking arrangements to 5,000 contenders. He then used computers to calculate the density of each arrangement. Doing so was more difficult than it sounds. The proof involved checking a series of mathematical inequalities using specially written computer code. In all, more than 100,000 inequalities were verified over a ten-year period.

Robert MacPherson, a mathematician at the Institute for Advanced Study in Princeton, New Jersey, and an editor of the *Annals*,

was intrigued when he heard about the proof. He wanted to ask Hales and his graduate student Sam Ferguson, who had assisted with the proof, to submit their finding for publication, but he was also uneasy about the computer-based nature of the work.

The *Annals* had, however, already accepted a shorter computer-aided proof — the paper, on a problem in topology, was published this March³. After sounding out his colleagues on the journal's editorial board, MacPherson asked Hales to submit his paper. Unusually, MacPherson assigned a dozen mathematicians to referee the proof — most journals tend to employ between one and three. The effort was led by Gábor Fejes Tóth of the Alfréd Rényi Institute of Mathematics in Budapest, Hungary, whose father, the mathematician László Fejes Tóth, had predicted in 1965 that computers would one day make a proof of Kepler's conjecture possible.

It was not enough for the referees to rerun Hales's code — they had to check whether the programs did the job that they were supposed to do. Inspecting all of the code and its inputs and outputs, which together take up three gigabytes of memory space, would have been impossible. So the referees limited themselves to consistency checks, a reconstruction of the thought processes behind each step of the proof, and then a

study of all of the assumptions and logic used to design the code. A series of seminars, which ran for full academic years, was organized to aid the effort.

But success remained elusive. Last July, Fejes Tóth reported that he and the other referees were 99% certain that the proof is sound. They found no errors or omissions, but felt that without checking every line of the code, they could not be absolutely certain that the proof is correct.

For a mathematical proof, this was not enough. After all, most mathematicians believe in the conjecture already — the proof is supposed to turn that belief into certainty. The history of Kepler's conjecture also gives reason for caution. In 1993, Wu-Yi Hsiang, then at the University of California, Berkeley, published a 100-page proof of the conjecture in the *International Journal of Mathematics*⁴. But shortly after publication, errors were found in parts of the proof. Although Hsiang stands by his paper, most mathematicians do not believe it is valid.

After the referees' reports had been considered, Hales says that he received the following letter from MacPherson: "The news from the referees is bad, from my perspective. They have not been able to certify the correctness of the proof, and will not be able to certify it in the future, because they have run out of energy ... One can speculate whether their process would have converged to a definitive answer had they had a more clear manuscript from the beginning, but this does not matter now."

Pyramid power:
Thomas Hales believes that computers will succeed where humans have failed in verifying his proof.



Star player: Johannes Kepler's conjecture has kept mathematicians guessing for 400 years.

The last sentence lets some irritation shine through. The proof that Hales delivered was by no means a polished piece. The 250-page manuscript consisted of five separate papers, each a sort of lab report that Hales and Ferguson filled out whenever the computer finished part of the proof. This unusual format made for difficult reading. To make matters worse, the notation and definitions also varied slightly between the papers.

Rough but ready

MacPherson had asked the authors to edit their manuscript. But Hales and Ferguson did not want to spend another year reworking their paper. "Tom could spend the rest of his career simplifying the proof," Ferguson said when they completed their paper. "That doesn't seem like an appropriate use of his time." Hales turned to other challenges, using traditional methods to solve the 2,000-year-old honeycomb conjecture, which states that of all conceivable tiles of equal area that can be used to cover a floor without leaving any gaps, hexagonal tiles have the shortest perimeter⁵. Ferguson left academia to take a job with the US Department of Defense.

Faced with exhausted referees, the editorial board of the *Annals* decided to publish the paper — but with a cautionary note. The paper will appear with an introduction by the editors stating that proofs of this type, which involve the use of computers to check a large number of mathematical statements, may be impossible to review in full. The matter might have ended there, but for Hales, having a note attached to his proof was not satisfactory.

This January, he launched the Flyspeck project, also known as the Formal Proof of Kepler. Rather than rely on human referees, Hales intends to use computers to verify

every step of his proof. The effort will require the collaboration of a core group of about ten volunteers, who will need to be qualified mathematicians and willing to donate the computer time on their machines. The team will write programs to deconstruct each step of the proof, line by line, into a set of axioms that are known to be correct. If every part of the code can be broken down into these axioms, the proof will finally be verified.

Those involved see the project as doing more than just validating Hales's proof. Sean McLaughlin, a graduate student at New York University, who studied under Hales and has used computer methods to solve other mathematical problems, has already volunteered. "It seems that checking computer-assisted proofs is almost impossible for humans," he says. "With luck, we will be able to show that problems of this size can be subjected to rigorous verification without the need for a referee process."

But not everyone shares McLaughlin's enthusiasm. Pierre Deligne, an algebraic geometer at the Institute for Advanced Study, is one of the many mathematicians who do not approve of computer-aided proofs. "I believe in a proof if I understand it," he says. For those who side with Deligne, using computers to remove human reviewers from the refereeing process is another step in the wrong direction.

Despite his reservations about the proof, MacPherson does not believe that mathematicians should cut themselves off from computers. Others go further. Freek Wiedijk, of the Catholic University of Nijmegen in the Netherlands, is a pioneer of the use of computers to verify proofs. He thinks that the process could become standard practice in mathematics. "People will look back at the turn of the twentieth century and say 'that is when it happened,'" Wiedijk says.


Whether or not computer-checking takes off, it is likely to be several years before Flyspeck produces a result. Hales and McLaughlin are the only confirmed participants, although others have expressed an interest. Hales estimates that the whole process, from crafting the code to running it, is likely to take 20 person-years of work. Only then will Kepler's conjecture become Kepler's theorem, and we will know for sure whether we have been stacking oranges correctly all these years.

George Szpiro writes for the Swiss newspapers NZZ and NZZ am Sonntag from Jerusalem, Israel. His book Kepler's Conjecture (Wiley, New York) was published in February.

1. Appel, K. & Haken, W. *Illinois J. Math.* **21**, 429–490 (1977).
2. Appel, K., Haken, W. & Koch, J. *Illinois J. Math.* **21**, 491–567 (1977).
3. Gabai, D., Meyerhoff, G. R. & Thurston, N. *Ann. Math.* **157**, 335–431 (2003).
4. Hsiang, W.-Y. *Int. J. Math.* **4**, 739–831 (1993).
5. Hales, T. C. *Discrete Comput. Geom.* **25**, 1–22 (2001).

Flyspeck
www.math.pitt.edu/~thales/flyspeck/index.html

19th C. MATHEMATICAL MODELS



Felix Klein's heritage

Considerable obstacles generally present themselves to the beginner, in studying the elements of Solid Geometry, from the practice which has hitherto uniformly prevailed in this country, of never submitting to the eye of the student, the figures on whose properties he is reasoning, but of drawing perspective representations of them upon a plane. . . .

I hope that I shall never be obliged to have recourse to a perspective drawing of any figure whose parts are not in the same plane.

Augustus de Morgan (1806–71).

- de Morgan, first President of the London Mathematical Society, was equally influential as an educator and a researcher.
- There is evidence that young children see more naturally in three than two dimensions.

(See discussion at www.colab.sfu.ca/ICIAM03/)



Coxeter's octahedral kaleidoscope (circa 1925)

Modern science is often driven by fads and fashion, and mathematics is no exception. Coxeter's style, I would say, is singularly unfashionable. He is guided, I think, almost completely by a profound sense of what is beautiful.

(Robert Moody)

In my book, Coxeter has been one of the most important 20th century mathematicians —not because he started a new perspective, but because he deepened and extended so beautifully an older esthetic. The classical goal of geometry is the exploration and enumeration of geometric configurations of all kinds, their symmetries and the constructions relating them to each other.

The goal is not especially to prove theorems but to discover these perfect objects and, in doing this, theorems are only a tool that imperfect humans need to reassure themselves that they have seen them correctly.


(David Mumford, 2003)

20th C. MATHEMATICAL MODELS



Ferguson's "Eight-Fold Way" sculpture

The Fergusons won the 2002 Communications Award, of the Joint Policy Board of Mathematics. The citation runs:



They have dazzled the mathematical community and a far wider public with exquisite sculptures embodying mathematical ideas, along with artful and accessible essays and lectures elucidating the mathematical concepts.

It has been known for some time that the *hyperbolic volume* V of the **figure-eight knot complement** is


$$\begin{aligned} V &= 2\sqrt{3} \sum_{n=1}^{\infty} \frac{1}{n \binom{2n}{n}} \sum_{k=n}^{2n-1} \frac{1}{k} \\ &= 2.029883212819307250042405108549 \dots \end{aligned}$$




Ferguson's "Figure-Eight Knot Complement" sculpture

In 1998, British physicist David Broadhurst conjectured $V/\sqrt{3}$ is a *rational linear combination* of

$$(2) \quad C_j = \sum_{n=0}^{\infty} \frac{(-1)^n}{27^n (6n + j)^2}.$$



Ferguson's
subtractive
image
of the
BBP Pi
formula



Indeed, as Broadhurst found, *using Ferguson's PSLQ*:

$$V = \frac{\sqrt{3}}{9} \sum_{n=0}^{\infty} \frac{(-1)^n}{27^n} \times \left\{ \frac{18}{(6n+1)^2} - \frac{18}{(6n+2)^2} - \frac{24}{(6n+3)^2} - \frac{6}{(6n+4)^2} + \frac{2}{(6n+5)^2} \right\}.$$

- Entering the following code in the *Mathematician's Toolkit*, at www.expmath.info:

```
v = 2 * sqrt[3] * sum[1/(n * binomial[2*n,n])
    * sum[1/k,{k, n,2*n-1}], {n, 1, infinity}]
```


```
pslq[v/sqrt[3],
    table[sum[(-1)^n/(27^n*(6*n+j)^2),
{n, 0, infinity}], {j, 1, 6}]]
```

recovers the solution vector


(9, -18, 18, 24, 6, -2, 0).

- The *first proof* that this formula holds is given in our new book.
- The formula is inscribed on each cast of the sculpture—marrying both sides of Helaman!

21st C. MATHEMATICAL MODELS



Knots 10_{161} (L) and 10_{162} (C) agree (R)*.



In NewMIC's Cave or Plato's?

*KnotPlot: from Little (1899) to Perko (1974) and on.



MORE of OUR 'METHODOLOGY'

1. (*High Precision*) computation of object(s).

2. *Pattern Recognition of Real Numbers*

$$\text{identify}(\sqrt{2}. + \sqrt{3}.) = \sqrt{2} + \sqrt{3}$$

(Inverse Calculator and 'identify')* or *Sequences* (Salvy & Zimmermann's 'gfun', Sloane and Plouffe's *Encyclopedia*).

3. Much use of 'Integer Relation Methods': †

✓ "Exclusion bounds" are especially useful.

✓ Great test bed for "Experimental Math".

4. Some automated theorem proving (Wilf-Zeilberger etc).


*ISC space limits: from 10Mb in 1985 to 10Gb today.

†*PSLQ, LLL, FFT*. Top Ten "Algorithm's for the Ages," Random Samples, Science, Feb. 4, 2000.

FOUR EXPERIMENTS

- 1. **Kantian** example: generating “the classical non-Euclidean geometries (hyperbolic, elliptic) by replacing Euclid’s axiom of parallels (or something equivalent to it) with alternative forms.”
- 2. The **Baconian** experiment is a contrived as opposed to a natural happening, it “is the consequence of ‘trying things out’ or even of merely messing about.”
- 3. **Aristotelian** demonstrations: “apply electrodes to a frog’s sciatic nerve, and lo, the leg kicks; always precede the presentation of the dog’s dinner with the ringing of a bell, and lo, the bell alone will soon make the dog dribble.”

- 4. The most important is **Galilean**: “a critical experiment – one that discriminates between possibilities and, in doing so, either gives us confidence in the view we are taking or makes us think it in need of correction.”
- It is also the only one of the four forms which will make Experimental Mathematics a serious enterprise.
- From Peter Medawar’s *Advice to a Young Scientist*, Harper (1979).




A Julia set


MILNOR

If I can give an abstract proof of something, I'm reasonably happy. But if I can get a concrete, computational proof and actually produce numbers I'm much happier.

I'm rather an addict of doing things on the computer, because that gives you an explicit criterion of what's going on. I have a visual way of thinking, and I'm happy if I can see a picture of what I'm working with.







ZEROES of 0 – 1 POLYNOMIALS



Data mining in polynomials


- The striations are unexplained!

WHAT YOU DRAW is WHAT YOU SEE



The price of metaphor is eternal vigilance (Arturo Rosenblueth & Norbert Wiener)

A NEW PROOF $\sqrt{2}$ is IRRATIONAL




- A *reductio* “proof without words”, published by Tom Apostol in the year 2000.
- But symbols are often more reliable than pictures.
- On to more detailed examples ...

TWO INTEGRALS

A. Why $\pi \neq \frac{22}{7}$:

$$0 < \int_0^1 \frac{(1-x)^4 x^4}{1+x^2} dx = \frac{22}{7} - \pi.$$

$$\left[\int_0^t \cdot = \frac{1}{7} t^7 - \frac{2}{3} t^6 + t^5 - \frac{4}{3} t^3 + 4t - 4 \arctan(t) . \right]$$





Archimedes: $\frac{223}{71} < \pi < \frac{22}{7}$

The Colour Calculator

B. *The sophomore's dream:*

$$\int_0^1 \frac{1}{x^x} dx = \sum_{n=1}^{\infty} \frac{1}{n^n}.$$

- Such have many implications for teaching — flagging issues of ‘Packing and unpacking’ concepts?



Pascal's Triangle modulo two

1, 1 1, 1 2 1, 1 3 3 1, 1 4 6 4 1, 1 5 10 10 5 1 ...

A. *A rational evaluation:*

$$\prod_{n=2}^{\infty} \frac{n^3 - 1}{n^3 + 1} = \frac{2}{3}.$$

...

B. *And a transcendent one:*

$$\prod_{n=2}^{\infty} \frac{n^2 - 1}{n^2 + 1} = \frac{\pi}{\sinh(\pi)}.$$

- The *Inverse Symbolic Calculator* can identify this product.
- \int, Σ, Π are now largely algorithmic not **black arts**.

HIGH PRECISION FRAUD

$$\sum_{n=1}^{\infty} \frac{[n \tanh(\pi)]}{10^n} \stackrel{?}{=} \frac{1}{81}$$

is valid to **268 places**; while

$$\sum_{n=1}^{\infty} \frac{[n \tanh(\frac{\pi}{2})]}{10^n} \stackrel{?}{=} \frac{1}{81}$$

is valid to just **12 places**.

- Both are actually transcendental numbers.

Correspondingly the *simple continued fractions* for $\tanh(\pi)$ and $\tanh(\frac{\pi}{2})$ are respectively

[0, 1, **267**, 4, 14, 1, 2, 1, 2, 2, 1, 2, 3, 8, 3, 1]

and

[0, 1, **11**, 14, 4, 1, 1, 1, 3, 1, 295, 4, 4, 1, 5, 17, 7]

- Bill Gosper describes how continued fractions let you “see” what a number is. “[I]t’s completely astounding ... **it looks like you are cheating God somehow.**”

CONVEX CONJUGATES and NMR(MRI)

The *Hoch and Stern information measure*, or *neg-entropy*, is defined in complex n -space by

$$H(z) = \sum_{j=1}^n h(z_j/b),$$

where h is convex and given (for scaling b) by:

$$h(z) \triangleq |z| \ln \left(|z| + \sqrt{1 + |z|^2} \right) - \sqrt{1 + |z|^2}$$

for quantum theoretic (NMR) reasons.

- Recall the *Fenchel-Legendre conjugate*

$$f^*(y) := \sup_x \langle y, x \rangle - f(x).$$

- Our *symbolic convex analysis* package (stored at www.cecm.sfu.ca/projects/CCA/) produced:

$$h^*(z) = \cosh(|z|).$$

- Compare the fundamental *Boltzmann-Shannon entropy*:

$$(z \ln z - z)^* = \exp(z).$$

★ I'd never have tried by hand!

- Knowing 'closed forms' helps:

$$(\exp \exp)^*(y) = y \ln(y) - y \{W(y) + W(y)^{-1}\}$$

where *Maple* or *Mathematica* recognize the complex *Lambert W* function

$$W(x)e^{W(x)} = x.$$

Thus, the conjugate's series is

$$-1 + (\ln(y) - 1)y - \frac{1}{2}y^2 + \frac{1}{3}y^3 - \frac{3}{8}y^4 + \frac{8}{15}y^5 + O(y^6)$$

SOME FOURIER INTEGRALS

Recall the *sinc* function

$$\operatorname{sinc}(x) := \frac{\sin(x)}{x}.$$

Consider, the seven highly oscillatory integrals below.*

$$I_1 := \int_0^\infty \operatorname{sinc}(x) dx = \frac{\pi}{2},$$

$$I_2 := \int_0^\infty \operatorname{sinc}(x) \operatorname{sinc}\left(\frac{x}{3}\right) dx = \frac{\pi}{2},$$

$$I_3 := \int_0^\infty \operatorname{sinc}(x) \operatorname{sinc}\left(\frac{x}{3}\right) \operatorname{sinc}\left(\frac{x}{5}\right) dx = \frac{\pi}{2},$$

...

$$I_6 := \int_0^\infty \operatorname{sinc}(x) \operatorname{sinc}\left(\frac{x}{3}\right) \cdots \operatorname{sinc}\left(\frac{x}{11}\right) dx = \frac{\pi}{2},$$

$$I_7 := \int_0^\infty \operatorname{sinc}(x) \operatorname{sinc}\left(\frac{x}{3}\right) \cdots \operatorname{sinc}\left(\frac{x}{13}\right) dx = \frac{\pi}{2}.$$

*These are hard to compute accurately numerically.

However,

$$\begin{aligned} I_8 &:= \int_0^\infty \operatorname{sinc}(x) \operatorname{sinc}\left(\frac{x}{3}\right) \cdots \operatorname{sinc}\left(\frac{x}{15}\right) dx \\ &= \frac{467807924713440738696537864469}{935615849440640907310521750000} \pi \\ &\approx 0.49999999999992646\pi. \end{aligned}$$

► When a researcher, using a well-known computer algebra package, checked this he – and the makers – concluded there was a “bug” in the software. Not so!

◇ Our analysis, via Parseval’s theorem, links the integral

$$I_N := \int_0^\infty \operatorname{sinc}(a_1 x) \operatorname{sinc}(a_2 x) \cdots \operatorname{sinc}(a_N x) dx$$

with the volume of the polyhedron P_N given by

$$P_N := \left\{ x : \left| \sum_{k=2}^N a_k x_k \right| \leq a_1, |x_k| \leq 1, 2 \leq k \leq N \right\}.$$

where $x := (x_2, x_3, \dots, x_N)$.


If we let

$C_N := \{(x_2, x_3, \dots, x_N) : -1 \leq x_k \leq 1, 2 \leq k \leq N\}$,
then

$$I_N = \frac{\pi \operatorname{Vol}(P_N)}{2a_1 \operatorname{Vol}(C_N)}.$$

► Thus, the value drops precisely when the constraint $\sum_{k=2}^N a_k x_k \leq a_1$ becomes *active* and bites the hypercube C_N . That occurs when $\sum_{k=2}^N a_k > a_1$.

In the above, $\frac{1}{3} + \frac{1}{5} + \dots + \frac{1}{13} < 1$, but on addition of the term $\frac{1}{15}$, the sum exceeds 1, the volume drops, and $I_N = \frac{\pi}{2}$ no longer holds.



- A somewhat cautionary example for too enthusiastically inferring patterns from seemingly compelling computation.

ENIAC: Integrator and Calculator

SIZE/WEIGHT: ENIAC had 18,000 vacuum tubes, 6,000 switches, 10,000 capacitors, 70,000 resistors, 1,500 relays, was 10 feet tall, occupied 1,800 square feet and weighed 30 tons.



SPEED/MEMORY: A 1.5GHz Pentium does 3 million adds/sec. ENIAC did 5,000 — 1,000 times faster than any earlier machine. The first stored-memory computer, ENIAC could store 200 digits.

ARCHITECTURE: Data flowed from one accumulator to the next, and after each accumulator finished a calculation, it communicated its results to the next in line.

The accumulators were connected to each other manually.

- The 1949 computation of π to 2,037 places took 70 hours.
- It would have taken roughly 100,000 ENIACs to store the Smithsonian's picture!

The computer has in turn changed the very nature of mathematical experience, suggesting for the first time that mathematics, like physics, may yet become an empirical discipline, a place where things are discovered because they are seen.

David Berlinski, "Ground Zero: A Review of *The Pleasures of Counting*, by T. W. Koerner," 1997.



A virtual fractal postcard

'PENTIUM FARMING' for BITS

B: Bailey, P. Borwein and Plouffe (1996) discovered a series for π (and other *polylogarithmic constants*) which allows one to compute hex-digits of π *without* computing prior digits.

► The algorithm needs very little memory and does not need multiple precision. The running time grows only slightly faster than linearly in the order of the digit being computed.

► The key, found by 'PSLQ' (below) is:

$$\pi = \sum_{k=0}^{\infty} \left(\frac{1}{16}\right)^k \left(\frac{4}{8k+1} - \frac{2}{8k+4} - \frac{1}{8k+5} - \frac{1}{8k+6}\right)$$

- Knowing an algorithm would follow they spent several months hunting for such a formula.
- Once found, easy to prove in *Mathematica*, *Maple* or by hand.

A most successful case of

**REVERSE
MATHEMATICAL
ENGINEERING**

► (Sept 97) Fabrice Bellard (INRIA) used a variant formula to compute 152 binary digits of π , starting at the *trillionth position* (10^{12}). This took 12 days on 20 work-stations working in parallel over the Internet.

► (August 98) Colin Percival (SFU, age 17) finished a similar ‘embarrassingly parallel’ computation of *five trillionth bit* (using 25 machines at about 10 times the speed). In *Hex*:

07E45733CC790B5B5979

The binary digits of π starting at the 40 trillionth place are

00000111110011111.

► (September 2000) The quadrillionth bit is '0' (used 250 cpu years on 1734 machines in 56 countries). From the 999,999,999,999,997th bit of π one has:

111000110001000010110101100000110

★ One of the largest computations ever!

Bailey and Crandall (2001) make a reasonable, hence very hard conjecture, about the **uniform distribution of a related chaotic dynamical system**. This conjecture implies:


Existence of a 'BBP' formula base b for α ensures the normality base b of α .

For $\log 2$ the dynamical system is

$$x_{n+1} \leftrightarrow 2\left(x_n + \frac{1}{n}\right) \pmod{1},$$

See www.sciencenews.org/20010901/bob9.asp .

A MISLEADING PICTURE



Polytopic except at one point?

The issue of paradigm choice can never be unequivocally settled by logic and experiment alone.

...

in these matters neither proof nor error is at issue. The transfer of allegiance from paradigm to paradigm is a conversion experience that cannot be forced.

- In *Who Got Einstein's Office?* ([Beurling](#))

And Max Planck, surveying his own career in his Scientific Autobiography, sadly remarked that 'a new scientific truth does not triumph by convincing its opponents and making them see the light, but rather because its opponents eventually die, and a new generation grows up that is familiar with it.'

A PARAPHRASE of HERSH

► Whatever the outcome of these developments, *mathematics is and will remain a uniquely human undertaking*. Indeed Reuben Hersh's arguments for a humanist philosophy of mathematics, as paraphrased below, become more convincing in our setting:

1. *Mathematics is human*. It is part of and fits into human culture. It does not match Frege's concept of an abstract, timeless, tenseless, objective reality.

2. *Mathematical knowledge is fallible*. As in science, mathematics can advance by making mistakes and then correcting or even re-correcting them. The "fallibilism" of mathematics is brilliantly argued in Lakatos' *Proofs and Refutations*.

3. *There are different versions of proof or rigor.* Standards of rigor can vary depending on time, place, and other things. The use of computers in formal proofs, exemplified by the computer-assisted proof of the **four color theorem** in 1977 (**1997**), is just one example of an emerging nontraditional standard of rigor.



4. *Empirical evidence, numerical experimentation and probabilistic proof all can help us decide what to believe in mathematics.* Aristotelian logic isn't necessarily always the best way of deciding.

5. *Mathematical objects are a special variety of a social-cultural-historical object.* Contrary to the assertions of certain post-modern detractors, mathematics cannot be dismissed as merely a new form of literature or religion. Nevertheless, many mathematical objects can be seen as shared ideas, like Moby Dick in literature, or the Immaculate Conception in religion.

► From “Fresh Breezes in the Philosophy of Mathematics”, *American Mathematical Monthly*, August-Sept 1995, 589–594.

► The recognition that “quasi-intuitive” analogies may be used to gain insight in mathematics can assist in the learning of mathematics.

And honest mathematicians will acknowledge their role in discovery as well. We should look forward to what the future will bring.

HILBERT

Moreover a mathematical problem should be difficult in order to entice us, yet not completely inaccessible, lest it mock our efforts. It should be to us a guidepost on the mazy path to hidden truths, and ultimately a reminder of our pleasure in the successful solution.

...

Besides it is an error to believe that rigor in the proof is the enemy of simplicity. (David Hilbert)

- In his '23' "Mathematische Probleme" lecture to the Paris International Congress, 1900 (see Ben Yandell's fine account in *The Honors Class*, AK Peters, 2002).

CHAITIN

I believe that elementary number theory and the rest of mathematics should be pursued more in the spirit of experimental science, and that you should be willing to adopt new principles. I believe that Euclid's statement that an axiom is a self-evident truth is a big mistake. The Schrödinger equation certainly isn't a self-evident truth! And the Riemann Hypothesis isn't self-evident either, but it's very useful. A physicist would say that there is ample experimental evidence for the Riemann Hypothesis and would go ahead and take it as a working assumption.*

*There is no evidence that Euclid ever made such a statement. However, the statement does have an undeniable emotional appeal.

CARATHÉODORY and CHRÉTIEN

I'll be glad if I have succeeded in impressing the idea that it is not only pleasant to read at times the works of the old mathematical authors, but this may occasionally be of use for the actual advancement of science.

- Constantin Carathéodory, at an MAA meeting in 1936 (retro-digital data-mining?).

A proof is a proof. What kind of a proof? It's a proof. A proof is a proof. And when you have a good proof, it's because it's proven.

(Jean Chrétien, 2002)


- The Prime Minister, explaining how Canada would determine if Iraq had WMDs, sounds a lot like Bertrand Russell!

GAUSS

► In Boris Stoicheff's often enthralling biography of Herzberg*, Gauss is recorded as writing:

It is not knowledge, but the act of learning, not possession but the act of getting there which generates the greatest satisfaction.

Fractal similarity in Gauss' discovery of modular functions



*Gerhard Herzberg (1903-99) fled Germany for Saskatchewan in 1935 and won the 1971 Chemistry Nobel.

A FEW CONCLUSIONS

- The traditional deductive accounting of Mathematics is a largely ahistorical caricature.*
- Mathematics is primarily about secure knowledge not proof, and the aesthetic is central.
- Proofs are often out of reach — understanding, even certainty, is not.
- Packages can make concepts accessible (Linear relations, Galois theory, Groebner bases).
- While progress is made “one funeral at a time” (Niels Bohr), “you can’t go home again” (Thomas Wolfe).

*Quotations are at jborwein/quotations.html

HOW NOT TO EXPERIMENT



Pooh Math

‘Guess and Check’
while

Aiming Too High

REFERENCES

1. J.M. Borwein, P.B. Borwein, R. Girgensohn and S. Parnes, “Making Sense of Experimental Mathematics,” *Mathematical Intelligencer*, **18**, (Fall 1996), 12–18.* [CECM 95:032]
2. Jonathan M. Borwein and Robert Corless, “Emerging Tools for Experimental Mathematics,” *MAA Monthly*, **106** (1999), 889–909. [CECM 98:110]
3. D.H. Bailey and J.M. Borwein, “Experimental Mathematics: Recent Developments and Future Outlook,” pp, 51-66 in Vol. I of *Mathematics Unlimited — 2001 and Beyond*, B. Engquist & W. Schmid (Eds.), Springer-Verlag, 2000. [CECM 99:143]

*All references are at www.cecm.sfu.ca/preprints.

4. J. Dongarra, F. Sullivan, “The top 10 algorithms,” *Computing in Science & Engineering*, **2** (2000), 22–23.
(See [personal/jborwein/algorithms.html](http://personal.jborwein/algorithms.html).)
5. J.M. Borwein and P.B. Borwein, “Challenges for Mathematical Computing,” *Computing in Science & Engineering*, **3** (2001), 48–53. [CECM 00:160].
6. J.M. Borwein and D.H. Bailey), **Mathematics by Experiment: Plausible Reasoning in the 21st Century**, and **Experimentation in Mathematics: Computational Paths to Discovery**, (with R. Girgensohn,) AK Peters Ltd, 2003-04.

► The web site is at www.expmath.info